

Medications Safety

In general, most medications should be avoided in pregnancy, *especially during the first trimester unless absolutely necessary*. However, some medications have a long history of safe use. If you are on medication, please discuss it with your provider. A few medications are dangerous in pregnancy, so be sure all your health care providers, including your dentist, know you are pregnant. The following medications can be used safely. If you have questions about the safety of additional medications, please discuss them with your provider or call us before taking them.

Safe Medications During Pregnancy

For Sleep	<ul style="list-style-type: none"> Benadryl (diphenhydramine), Tylenol PM, Unisom (doxylamine), Melatonin
For Fever, Pain, Headache, or Muscle Soreness	<p>Note: DO NOT take aspirin or ibuprofen unless directed by your provider.</p> <ul style="list-style-type: none"> Tylenol (acetaminophen) - regular or extra-strength
For Morning Sickness	<ul style="list-style-type: none"> Vitamin B6 (50mg one-three times a day) - works best when also taken with Unisom (1/2 tablet once or twice a day, can be sedating), Emetrol, Benadryl (diphenhydramine) (25-50mg), motion sickness medication such as Dramamine
For Nausea, Upset Stomach, or Gas	<ul style="list-style-type: none"> Emetrol, Mylanta (aluminum hydroxide), Gas X (simethicone)
For Heartburn, Acid Reflux, and Indigestion	<p>Note: Non-medication options include eating frequent, smaller meals, avoid laying down for 1 hour after eating, and avoiding spicy or acidic foods.</p> <ul style="list-style-type: none"> Tums (calcium carbonate) (no more than 1,000mg/day), Pepcid AC (famotidine), Mylanta, Maalox
For Constipation	<p>Note: Non-medication options include increasing water and vegetable intake.</p> <ul style="list-style-type: none"> (In moderation) Fiber source such as Metamucil or Citrucel, stool softener such as Colace (docusate), glycerin suppository, Milk of Magnesia, Senokot, Miralax
For Hemorrhoids	<p>Note: Non-medication options include increasing your water intake.</p> <ul style="list-style-type: none"> Fiber supplements, Tucks pads, witch hazel compress, Anusol, Preparation H
For Cold or Other Upper Respiratory Illnesses	<p>Note: Avoid pseudoephedrine in the first trimester of pregnancy. If necessary after the first trimester, limit to 2-3 days.</p> <ul style="list-style-type: none"> Nasal saline spray, nasal steroid spray, Benadryl (diphenhydramine), Afrin (oxymetazoline hydrochloride), Zyrtec (cetirizine), Allegra (fexofenadine), Claritin (loratadine), Sudafed (pseudoephedrine), Tylenol Cold and Sinus (acetaminophen and pseudoephedrine)
For Cough	<ul style="list-style-type: none"> Cough drops, Robitussin DM (dextromethorphan and pseudoephedrine), Dimetapp (brompheniramine and pseudoephedrine)
For Sore Throat	<p>Note: Non-medication options include gargling with salt water.</p> <ul style="list-style-type: none"> Chloraseptic throat spray, Tylenol (acetaminophen), Luden's throat drops
For Diarrhea	<ul style="list-style-type: none"> Imodium AD (loperamide)
For Yeast Infection	<ul style="list-style-type: none"> Monistat or other vaginal yeast creams
For Allergies	<ul style="list-style-type: none"> Benadryl (diphenhydramine), Claritin (loratadine), Zyrtec (cetirizine)

The above over-the-counter remedies are to be used for temporary ailments. If your condition persists, please call us.